



phexxi™

(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

pHinally!

A NEW, FDA-APPROVED,  
**HORMONE-FREE**  
BIRTH CONTROL



**\$0 CO-PAY FOR YOUR  
1<sup>ST</sup> PRESCRIPTION**

See last page for details.

Please see [Prescribing Information](#), including  
[Patient Information](#).

**phexxi™**  
(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

Are you one of the millions of women who  
have said **enough is enough** to hormones?

NOW, THERE'S **PHEXXI™**!

## WHAT IS **PHEXXI™**?

Phexxi™ is a prescription vaginal gel used to prevent pregnancy in females who choose to use an on-demand method of birth control. Phexxi™ is only effective when used immediately **before** (or up to one hour before) each act of vaginal sex. It is not effective when used after vaginal sex.



## HOW **PHEXXI™** WORKS

Vaginas have a pH level that is inhospitable to sperm. When semen enters the vagina, the pH level is increased to help sperm survive. Phexxi™ works to maintain vaginal pH, which reduces the sperm's mobility, lowering the chance of sperm reaching the egg.

## **IMPORTANT SAFETY INFORMATION**

### **WHAT ARE THE POSSIBLE SIDE EFFECTS OF PHEXXI™?**

There have been a few cases of urinary bladder infection and kidney infection reported in clinical studies. One of the cases was serious. Avoid Phexxi™ if you have had repeated urinary tract infections or other urinary tract problems.

Please see [Prescribing Information](#), including [Patient Information](#).

**phexxi™**  
(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

## **PHEXXI™ IS QUICK AND EASY TO USE**

Phexxi™ is designed to be used in the moment  
and works immediately upon application.



Not actual size.

## **PHEXXI™ PUTS YOU IN CONTROL**

- Insertion of the Phexxi™ applicator is similar to inserting a tampon
- Phexxi™ doesn't have to be used alone. You can use it with condoms or diaphragms, but not with a vaginal ring
- For your convenience, Phexxi™ comes in individually wrapped, single-dose applicators
- Phexxi™ is effective up to 1 hour upon application. After 1 hour, an additional dose of Phexxi™ needs to be used before each act of sex

Phexxi™ is not effective when used after vaginal sex.

### **IMPORTANT SAFETY INFORMATION (CONT'D)**

#### **WHAT ELSE SHOULD I KNOW ABOUT USING PHEXXI™?**

Phexxi™ does not protect against any sexually transmitted diseases, including HIV. Avoid using Phexxi™ with a vaginal ring.

Please see [Prescribing Information](#), including [Patient Information](#).

**phexxi™**  
(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

*Phexxi™ is FDA-approved*

**PHEXXI™** HAS  
DEMONSTRATED THAT  
IT IS EFFECTIVE AT  
PREVENTING PREGNANCY

Phexxi™ has not been associated with side effects  
such as blood clots or stroke.

### POSSIBLE SIDE EFFECTS OF **PHEXXI™**

There have been a few cases of urinary bladder infection and kidney infection reported in clinical studies. One of the cases was serious. Avoid Phexxi™ if you have repeated urinary tract infections or other urinary tract problems.

---

See additional safety information  
throughout the brochure.

## **IMPORTANT SAFETY INFORMATION (CONT'D)**

### **WHAT ARE THE POSSIBLE SIDE EFFECTS OF PHEXXI™? (CONT'D)**

The **most common side effects** were vaginal burning, vaginal itching, vaginal yeast infection, urinary tract infection, vaginal area discomfort, bacterial vaginosis, and vaginal discharge. Women also reported genital discomfort, pain while urinating, and vaginal pain. Some male partners reported genital discomfort.

Please see [Prescribing Information](#), including [Patient Information](#).

**pH**ABULOUS!

Ask your healthcare provider if **Phexxi**<sup>™</sup> is right for you.

**phexxi**<sup>™</sup>  
(lactic acid, citric acid, and potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

**pH**ANTASTIC!

SAVE WITH THE PHEXXI<sup>™</sup> CO-PAY CARD

Under the Affordable Care Act, Phexxi<sup>™</sup> may be available for \$0 co-pay, co-insurance, or deductible.



[Click here](#)  
to access the  
co-pay card.

Learn more at [phexxi.com](http://phexxi.com).

## IMPORTANT SAFETY INFORMATION (CONT'D)

### WHAT ELSE SHOULD I KNOW ABOUT USING PHEXXI<sup>™</sup>? (CONT'D)

Contact your healthcare provider if you are experiencing severe genital irritation or discomfort or urinary tract symptoms. Avoid Phexxi<sup>™</sup> if you or your sexual partner is allergic to lactic acid, citric acid, potassium bitartrate, or any of the ingredients in Phexxi<sup>™</sup>. Stop using Phexxi<sup>™</sup> if you develop an allergic reaction.

**Please report side effects by contacting Evofem Biosciences<sup>®</sup> toll-free at 1-833-EVFM BIO or contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

Please see [Prescribing Information](#), including [Patient Information](#).

**EVOFEM**  
BIOSCIENCES<sup>®</sup>

Trademarks are owned by Evofem Biosciences.  
© 2020 Evofem Biosciences.  
EVFM-US-000634 • September 2020